

Eco-friendly Synthesis Of 2-phenylchroman-4-one by Using Ionic Liquid

Dr. Arshia Parveen, Department of Chemistry, B. Raghunath College, Parbhani-431401 (MS).

Introduction

Flavanones are a group of common and naturally occurring polyphenolic compounds that are widely found in the plant kingdom (Geissman et al, 1969). They occur naturally as plant pigments in a broad range of fruits and vegetables as well as beverages such as tea, red wine, coffee (Murray et al 1996). Flavanone have been reported to exert multiple biological effects including antimicrobial (Proestos et al, 2005), cytotoxicity (Yenjai, 2004), anti-inflammatory (Furuta et al, 2004) as well as anti-tumor activities (Xia et al, 2000). In this regard, several flavanones bearing hydroxyl groups on the A or B ring have been reported to be potential antioxidant agents. It is now well established that such potency is mainly due to the ability of hydroxyl groups to donate hydrogen which enable the flavanone to undergo a redox reaction that helps them to scavenge free radicals (Hertog et al, 1993). In addition, the presence of hydroxyl groups in the skeleton also contribute to high affinity for proteins and therefore acts as inhibitors of microbial enzymes (Prusky et al, 1993) and inhibition of NADH dehydrogenase of mitochondrial inner membranes (Ravanel et al, 1989). Prenylated flavanones are a unique class of naturally occurring flavanone characterized by the presence of a prenylated side chain in the flavonoid skeleton. It was reported that one phenolic group and certain degree of lipophilicity are required for the activity of the flavonones (Laks et al, 1989). Substitution of the flavanoid ring system with prenyl groups would increase their lipophilicity and consequently enhance their interaction with cellular membranes (Baron et al, 1989). Different methods are used for the synthesis of flavones, includes Allan-Robinson synthesis (Banerji et al, 1980), synthesis from chalcones (Hoshino et al, 1986) and via intramolecular Wittig reaction (LeFloch et al, 1986). The most common method used involves Baker-Venkattram arrangement. In this method 2-hydroxyacetophenone are converted to benzoyl ester, which in presence of base (pyridine/KOH) form 1,3-diketones. The diketones are further cyclized under strong acidic condition to afford the flavones (Balogh et al, 1993, Chisen et al, 1997). In recent development of such dehydrative cyclization it includes the use of Amberlyst 15 (Hoshino et al, 1987), Co^{III} (sulpr)OH (Nishinaga et al, 1982), FeCl_3 (Zubaidha et al, 2005), $\text{Br}_2/\text{CHCl}_3$ (Garg et al, 1994), EtOH/HCl (Jung et al, 2001), clay (Verma et al, 1998), NaOAc/AcOH (Kumar et al, 1999) and H_2SO_4 under microwave irradiation (Tsukayama et al, 2003). Prenylated flavanone is a unique class of naturally occurring flavonoids characterized by the presence of a prenylated side chain in the flavonoid skeleton. It was reported that one phenolic group and certain degree of lipophilicity are required for the activity of the flavonoids (Baron et al, 1989). Substitution of the flavonoid ring system with prenyl groups would increase their lipophilicity and consequently enhance their interaction with cellular membranes. 4',5,7-Trihydroxy-3'-prenylflavanone (1) has been isolated for the first time in 1989 from the chloroform extract of the stem bark of *Erythrinaeriothiocha* (Nkengfack et al, 1989). The chemical and pharmaceutical industries are always under the pressure to find out environmental friendly organic reaction methodologies. Microwave irradiation is used for a variety of organic reactions due to their use in a rapid and cleaner synthesis of organic compounds (Varma et al, 1993 and Marrero-Terrero et al, 1996 and Benalloum et al, 1998 and Lerestif et al, 1997).

Ionic liquids are possible green catalyst acts as alternatives for several catalytic reactions. Ionic liquids attracted attention of researchers due to their mild reaction conditions, short reaction times and better yield, solvating ability and easy recyclability (Welton et al, 1999 and Wassercheid et


al, 2000 and Sheldon et al, 2001 and Zhao et al, 2002). Various reactions have been reported recently using ionic liquids as a catalyst, reaction media (Rajgopal et al, 2003 and Jarikoteet al, 2003 and Gholap et al, 2004 and Panchgalle, 2004) and as rate enhancers (Madjeet al, 2004). (Sarda et al, 2006) synthesized a high yielding and fast method for smooth conversion of substituted 1-(2-hydroxy phenyl)-3-phenyl-1,3-propane diones to corresponding chromen-4-one under microwave irradiations using ionic liquid [EtNH₃]⁺NO₃⁻ is firstly reported.

Experimental

The α - β -unsaturated carbonyl compounds **1** (1 mmol) was added in ionic liquid EAN (2 mmol) and irradiated into a domestic microwave oven at 250^oC for 20-25 sec. The reaction was monitored on TLC. After completion of the reaction, the mixture was extracted 5 X 20 ml. of ethyl acetate: petroleum ether (50%+50%). Compound comes in organic layer, was again treated with water, brine & dried over MgSO₄. Organic solvent is evaporated to afford pure flavanones **2**. Further, ionic liquid was dried under reduced pressure & reused for another reaction gives same yield. The recovery percentage of ionic liquid is satisfactory. The obtained products **2a**-**2p** were identified by comparison with authentic samples ¹H NMR and their melting points.

Results and discussion


From the observations of the literature it is noted that the yield is low to moderate in the conventional methods some time due to catalyst or due to solvent therefore we choose a novel methodology in which we used IL as catalyst as well as solvent so it will become green synthesis. (Scheme-1) We used EAN ethyl ammonium nitrate.



Scheme-1 synthesis of flavanone using EAN ionic liquid

Herein we wish to report for the first time a novel synthesis of flavanones **2** promoted by ionic liquid catalyst, ethyl ammonium nitrate [EtNH₃]⁺NO₃⁻ under microwave irradiation in excellent yield with shorter reaction time (Scheme 1). The ionic liquid can be recycled and reused several times. The ionic liquid [EtNH₃]⁺NO₃⁻ was prepared as per literature method³².

The effect of microwave heating was studied considering different solid supports and catalyst. After some experimentation the best procedure for the irradiation is described as follows. The reaction mixtures were heated successively for 10 sec periods followed by a 5 sec cooling interval between irradiations. The method was designed to avoid over heating of reactants. Since the unmodified house hold microwave oven lacks the special attributes of commercial reactors in terms of power and temperature control. In addition, because a slandered house hold microwave oven was used, a experiment to test the homogeneity of the irradiating field was conducted. In a typical reaction, the 2'-hydroxychalcone (**1**) in ionic liquid [EtNH₃]⁺NO₃⁻ was irradiated under domestic microwave oven in a specified time. The progress of the reaction was monitor by TLC. After completion of reaction, aqueous work up afforded pure flavanones (**2**) in 85% yield. To evaluate the synthetic utility of the process, various substituted chalcones were prepared by the previous (scheme1) and subjected to the reaction under microwave irradiation. The results are shown in Table-1.



Scheme 2 plausible mechanism for the synthesis of flavanone **2**

ANTIMICROBIAL ACTIVITY

The antibacterial activities of the synthesized compounds (d) and (f) were studied against four bacteria, viz. *Bacillus subtilis* (G+), *Escherichia coli* (G-), *Staphylococcus aureus* (G+) and *Pseudomonas aeruginosa* (G-). For the detection of antibacterial activities, the filter paper discs diffusion method was used (Tsukayama et al, 2003). Streptomycin sulphate was used as positive control. Nutrient agar (NA) was used as basal medium for test bacteria. The discs were prepared by impregnating them in methanol solution of each sample (1 mg/1 mL).




Table 1 Synthesis of Flavonone **2** by using EAN ionic liquid in microwave oven.

Pdt	R1	R2	R3	R4	R5	R6	R7	Time sec	^a Y%
2a	H	H	H	H	H	H	H	10	89
2b	H	H	H	H	Cl	H	H	12	90
2c	H	H	H	H	OMe	H	H	14	87
2d	H	H	Me	H	OMe	H	H	16	86
2e	H	H	H	H	OMe	OMe	H	12	92
2f	H	OH	H	H	H	H	H	15	93
2g	H	H	H	H	H	H	cl	17	95
2h	H	H	H	H	H	OMe	H	13	89
2i	H	H	H	H	Br	H	H	10	93
2j	H	H	H	H	H	H	NO ₂	12	90
2k	H	H	H	H	H	H	OMe	11	90
2l	H	H	H	H	H	H	Br	13	89
2m	H	H	H	H	OC ₆ H ₁₃	H	H	14	91
2n	H	H	H	H	OC ₉ H ₁₇	H	H	17	92
2o	H	OMe	H	Me	Br	H	H	18	90
2p	H	OC ₄ H ₉	H	H	OC ₄ H ₉	H	H	20	89

^a isolated yield

Each culture was prepared to a turbidity equivalent to McFarland and spread on the test tube. The paper disc containing the compound were placed on the agar surface previously inoculated with suspension of each microbes to be tested. All determinations were made in duplicate.

Inhibition diameter were determined after incubation at $37^{\circ}\text{C} \pm 1$ for 24 h. The antimicrobial activity was indicated by the presence of the clear inhibition zones around each disc.

Minimum inhibition concentration :

The determination of the minimum inhibitory concentration (MIC), the serial dilution technique was followed using nutrient broth medium. The MIC was defined as the lowest concentration of samples that had restricted the growth of microbial. The MIC value of compound (c) were determined against *Escherichia coli* (G-).

Conclusions

In summary we have demonstrated an efficient and mild protocol for the synthesis of flavanones by cyclization of chalcones using EAN under microwave irradiation and dehydrative cyclization of 1,3-(diaryl) diketones to flavanones in presence of ionic liquids. Shorter reaction time, simple reaction conditions and higher yield render this microwave irradiation method superior. The method is clean and simple, which can be used as an alternative to the existing methods.

ANTIMICROBIAL SCREENING

The antibacterial activity of compounds (d) and (f) has been assayed at the concentration 1000 $\mu\text{g/mL}$ against four human pathogenic bacteria. Among them two were gram-positive and the other two were gram negative. The inhibitory effect of compounds (d) and (f) against these organisms are given in table 2. The screening results indicate that only compound (f) was active against a gram-negative bacteria, *Escherichia coli* with a mean zone of inhibition 12.5 ± 0.3 mm (table 2).

Determination of the minimum inhibitory concentration (MIC) :

The active sample in the disc diffusion method was then tested for its activity by the serial dilution method to determine the minimum inhibition concentration (MIC-value). The MIC value obtained for flavanone (f) was 1000 $\mu\text{g/mL}$ against *Escherichia coli*.

Table 2. Antibacterial screening for the compounds (b) and (g)

Organism	chalcones	flavanone	Streptomycin sulphate
Bacillus subtilis	–	–	22.0 ± 0.3
Staphylococcus aureus	–	–	22.5 ± 0.7
Escherichia coli	–	12.5 ± 0.3	22.0 ± 0.0
Pseudomonas aeruginosa	–	–	22.0 ± 0.0

Diameter of the zone of inhibition (mm)

References:

1. Geissman T A ,Crout D H G .Organic chemistry of secondary plantmetabolites(California: Freeman, Cooper and Company) pp. **1969** ;183–230.
2. Murray M T . Encyclopedia of nutritional supplements (California: Prima Publishing) pp. **1996**; 320–331.

3. Proestos C, Boziaris I.S, Nychas JE. Food Chem. **2005**; (93) :1998.
4. Yenjai C, Prasanphen K, Daodee S , Kittakoop P. Fitoterapia**2004** ;(75):89.
5. Furuta T, Kimura, T, Kondo S, Wakimoto T, Nukaya H, Tsuji K, Tanaka K. Tetrahedron **2004**;(60): 9375.
6. Xia Y, Yang ZY, Xia. P, Bastow K F, Nakanishi Y, Lee K H. Bioorg. Med. Chem. Lett.**2000**;(10): 699.
7. Hertog MG L. Lancet **1993** ;(342) :1007.
8. Prusky D, Keen N T. Plant Dis. **1993** ;(77): 114.
9. Ravanel P, Creuzet S, Tissut M. Phytochemistry **1989**;(27): 87.
10. Laks P E, Pruner M S. Phytochemistry**1989**;(28): 87.
11. Baron D, Ragai K I. Phytochemistry**1989**;(27) :87.
12. Banerji A, Goomer N, *Synthesis***1980**, 874.
13. Hoshino Y, Oohinata T, Takeno N. Bull. Chem. Soc. Jpn**1986**;(59): 2351.
14. LeFloch'h Y, LeFeuvre M. Tetrahedron Lett**1986**; (27): 2751.
15. Balogh M, Laszlo P. Organic Chemistry Using Clays, Springer, Berlin,**1993**; Chisen J, Chisen IC, Rafelt JS, Macquarrie DJ , Clark JH. Chem. Commun**1997**; 2203.
16. Hoshino Y ,Takino N. Bull. Chem. Soc. Jpn. **1987**;(60):1919-1920.
17. Nishinaga A, Ando H, Maruyama K, Mashino T. *Synthesis***1982**; 839.
18. Zubaidha PK, Hashmi AM ,Bhosale RS. HetrocyclicCommun. **2005**;(11): 9100.
19. Garg S, Ishar M P S, Sarin R, R.P. Gandhi, Indian J. Chem. Soc. **1994**; (33B):1123-1128.
20. Jung J C, Min J P , Park O S. Synth. Commun. **2001**; 31(12):1837-21.
21. Verma R S, Saini R K, Kumar D, J. Chem. Res. (S). **1998**; 348- 349.
22. Kumar P E, Prashad K J R, Indian J. Chem. **1999**; (38B): 1277-1279.
23. Tsukayama M, Kawamura Y, Ishizuka T, Hayas S, F. Torii F.Heterocycles. **2003**; 60 (12), 2775.
24. Baron D ,Ragai K I. Phytochemistry**1989**; (27)87.
25. Nkengfack A E, Sanson D R , Tempesta M S J. Nat. Prod. **1989**;(52) 320.
26. Varma R S, Varma M , Chatterjee A K.J. Chem. Soc., PerkinTrans.**1993**1,;999. Varma R S, Chatterjee A K, M. Varma M. Tetrahedron Lett. **1993**;(34): 4603.
27. Lerestif J M, Toupet L, Sinbandhit S, Tonnerd F, Bazureaau J P, HanelinJ, Terahedron**1997**;(53):6351. Marrero-Terrero AL, Loupy A. Synlett.**1996**;245. Benalloum A, LabiadB , VilleminD. Chem. Commun. **1998**; 386.
28. Welton T. Chem. Rev.**1999**;(99): 2071. Wassercheid P, Keim W, Angew.Chem. Int. Ed. **2000**;(39): 3772. Sheldon R. Chem. Commun.**2001**; 2399. Zhao D, Wu M, Kou Y, Min K. Catal. Today. **2002**; (1): 2654.
29. RajgopalR, Jarikote DV, LahotiRJ, ThomasD, Srinivasan KV. *Tetrahedron Lett.* **2003**; (44):1615. Jarikote D V, Siddiqui S A, Rajgopal R., Thomas D, Lahoti R J , Srinivasan K V. Tetrahedron Lett. **2003**; (44):1835. Gholap A R Venkatesan K, Thomas D, Lahoti R J, Srinivasan K V, Green Chem. **2004**; (6),:147-150. Panchgalle S P, Kalkote U R, Nipahadkar P S, Joshi P N, Chavan S P, Chaphekar G M, Green Chem. **2004**; (6): 308-309.
30. Madje B R, Shindalkar S S , Shingare M S. Indian Journal of Hetrocy. Chem. **2004**; (14):87-88.
31. Sarda S R, Pathan M K, Paike W, Pechmase P R, Jadhav W N, Pawar R P . *Arkivoc***2006**;(16):43-80.